

### Quiz 4

A record player is set for  $33 \frac{1}{3}$  revolutions per minute (rpm). It is found that a penny will stick with the platter as long as its distance from the platter center is less than 15 cm. (A) Draw a free body diagram of the penny sitting on the rotating platter. Show and name all forces acting on the penny. Show the direction of the acceleration (if there is any). (B) Calculate the coefficient of static friction for the penny on the platter.

$$\omega = 33 \frac{1}{3} \frac{\text{rev}}{\text{min}} \cdot \frac{1 \text{ min}}{60 \text{ s}} \cdot \frac{2\pi \text{ rad}}{1 \text{ rev}} = 3.49 \frac{\text{rad}}{\text{sec}}$$

$$N = mg$$

$$f = \mu_s N$$

usually  $\leq$  but in this problem at limit


$$= ma$$

$$\mu_s mg = m \omega^2 r \Rightarrow \mu_s = \frac{\omega^2 r}{g}$$

$$= \frac{(3.49)^2 (0.15)}{9.8}$$


$$= .187$$

↑ unitless



### Quiz 4

A record player is set for  $33 \frac{1}{3}$  revolutions per minute (rpm). It is found that a penny will stick with the platter as long as its distance from the platter center is less than 15 cm. (A) Draw a free body diagram of the penny sitting on the rotating platter. Show and name all forces acting on the penny. Show the direction of the acceleration (if there is any). (B) Calculate the coefficient of static friction for the penny on the platter.



### Quiz 5

Stuntman Rupert has decided his Terminator+bike (mass = 100 kg) should have a horizontal velocity of 14 m/s for the crack-jump scene of Godzilla vs. Terminator. Since he's not really the Terminator, he could not achieve the required speed using his own muscles. So he's decided to build a frictionless ramp out of view of the camera to provide all the required energy. However, the movie frame will include 3 m on the rough surface before the crack (which has a frictional force of 800 N). (A) Calculate the work done by the frictional force as the bike travels over the rough surface. (B) Calculate the bike+rider kinetic energy (at the crack edge) required for a successful jump. (C) Calculate the height  $h$  at which Rupert should start.

A:  $W = Fd \cos \theta = Fd (-1) = -800 \cdot 3 = -2400 \text{ J}$


B  $KE = \frac{1}{2} mv^2 = \frac{1}{2} \cdot 100 \cdot 14^2 = 9.8 \times 10^3 \text{ J}$

$$W_{nc} = (KE_f + PE_f) - (KE_i + PE_i)$$

$\uparrow$                      $\uparrow$                      $\uparrow$                      $\uparrow$   
 $-2400$              $9.8 \times 10^3$      $0$                      $0$                      $mgh$


$$mgh = +2400 + 9.8 \times 10^3$$

$$h = \frac{(2400 + 9800)}{100 \cdot 9.8} = 12.5 \text{ m}$$



### Quiz 5

Stuntman Rupert has decided his Terminator+bike (mass = 100 kg) should have a horizontal velocity of 14 m/s for the crack-jump scene of Godzilla vs. Terminator. Since he's not really the Terminator, he could not achieve the required speed using his own muscles. So he's decided to build a frictionless ramp out of view of the camera to provide all the required energy. However, the movie frame will include 3 m on the rough surface before the crack (which has a frictional force of 800 N). (A) Calculate the work done by the frictional force as the bike travels over the rough surface. (B) Calculate the bike+rider kinetic energy (at the crack edge) required for a successful jump. (C) Calculate the height  $h$  at which Rupert should start.



## Quiz 6

A 5.5 kg bowling ball moving 9 m/s makes an elastic, head-on collision with a .85 kg bowling pin which moves straight ahead (i.e., everything remains in a line). Find the post collision velocities of the bowling ball and bowling pin.

$m_1 \leftarrow v_1$        $m_2 \leftarrow v_2 = 0$   
 $v_1'$        $v_2'$

$m_1 v_1 + m_2 v_2 = m_1 v_1' + m_2 v_2'$       momentum  
 $m_1 v_1 = m_1 v_1' + m_2 v_2'$

$m_1 v_1 = m_1 v_1' + m_2 (v_1 + v_1')$

$(m_1 - m_2) v_1 = (m_1 + m_2) v_1'$

$\frac{(m_1 - m_2)}{(m_1 + m_2)} v_1 = v_1'$

$\frac{5.5 - .85}{5.5 + .85} 9 = v_1'$

$6.59 \text{ m/s} = v_1'$

KE conservation is equivalent to  
 $(v_1 - v_2) = -(v_1' - v_2')$

$v_1 = v_2' - v_1'$   
 $v_1 + v_1' = v_2'$

$9 + 6.59 = v_2'$   
 $15.6 \text{ m/s} = v_2'$

## Quiz 6

A 5.5 kg bowling ball moving 9 m/s makes an elastic, head-on collision with a .85 kg bowling pin which moves straight ahead (i.e., everything remains in a line). Find the post collision velocities of the bowling ball and bowling pin.