

# Fields and Moments of a Moving Electric Dipole

V. Hnizdo<sup>1</sup>

National Institute for Occupational Safety and Health, Morgantown, WV 26505

Kirk T. McDonald

Joseph Henry Laboratories, Princeton University, Princeton, NJ 08544

(November 29, 2011)

## 1 Problem

Discuss the fields of a moving point electric dipole, and characterize these fields in terms of multipole moments. It suffices to consider uniform motion with velocity  $v$  small compared to the speed  $c$  of light in vacuum.

## 2 Solution

*This note is an elaboration of [1].*

The relativistic transformations of densities  $\mathbf{P}$  and  $\mathbf{M}$  of electric and magnetic dipole moments (also called polarization and magnetization densities, respectively) were first discussed by Lorentz [2], who noted that they follow the same transformations as do the magnetic and electric fields  $\mathbf{B} = \mathbf{H} + 4\pi\mathbf{M}$  and  $\mathbf{E} = \mathbf{D} - 4\pi\mathbf{P}$ , respectively,

$$\mathbf{P} = \gamma \left( \mathbf{P}_0 + \frac{\mathbf{v}}{c} \times \mathbf{M}_0 \right) - (\gamma - 1) \frac{(\mathbf{v} \cdot \mathbf{P}_0) \mathbf{v}}{c^2}, \quad \mathbf{M} = \gamma \left( \mathbf{M}_0 - \frac{\mathbf{v}}{c} \times \mathbf{P}_0 \right) - (\gamma - 1) \frac{(\mathbf{v} \cdot \mathbf{M}_0) \mathbf{v}}{c^2}, \quad (1)$$

where the inertial lab frame moves with velocity  $\mathbf{v}$  with respect to the (inertial) rest frame of the polarization densities, and  $\gamma = 1/\sqrt{1 - v^2/c^2}$ .

Considerations of the fields moving electric dipoles perhaps first arose in the context of Čerenkov radiation when the dipole velocity  $v$  exceeds the speed of light  $c/n$  in a medium of index of refraction  $n$  [3]. Later discussions by Frank of his pioneering work are given in [4, 5].

As noted by Frank [5], specialization of eq. (1) to point electric and magnetic dipole moments  $\mathbf{p}$  and  $\mathbf{m}$ , and to low velocities, leads to the forms<sup>2</sup>

$$\mathbf{p} = \mathbf{p}_0 + \frac{\mathbf{v}}{c} \times \mathbf{m}_0, \quad \mathbf{m} = \mathbf{m}_0 - \frac{\mathbf{v}}{c} \times \mathbf{p}_0, \quad \mathbf{p}_0 = \mathbf{p} - \frac{\mathbf{v}}{c} \times \mathbf{m}, \quad \mathbf{m}_0 = \mathbf{m} + \frac{\mathbf{v}}{c} \times \mathbf{p}, \quad (2)$$

where  $\mathbf{p}_0$  and  $\mathbf{m}_0$  are the moments in the rest frame of the point particle, while  $\mathbf{p}$  and  $\mathbf{m}$  are the moments when the particle has velocity  $\mathbf{v}$  in the lab frame. However, the fields associated with an electric and/or magnetic dipole moving at low velocity are not simply the instantaneous fields of the moments  $\mathbf{p}$  and  $\mathbf{m}$ , which leads to ambiguities in interpretations of the fields as due to moments.

Before deducing the fields, we note a past misunderstanding regarding eq. (2).

---

<sup>1</sup>This note was written by the author in his private capacity. No official support or endorsement by the Center for Disease Control and Prevention is intended or should be inferred.

<sup>2</sup>Ref. [5] recounts a past controversy that these transformations might involve the index  $n$  if the magnetic dipole were not equivalent to an Ampèrian current loop.

## 2.1 Fisher's Claim

In sec. III of [6] it is claimed that the magnetic moment of a point electric dipole  $\mathbf{p}_0$  which moves with velocity  $\mathbf{v}$  can be calculated according to

$$\mathbf{m} = \frac{1}{2c} \int \mathbf{r} \times \mathbf{J} d\text{Vol} = \frac{1}{2c} \int \mathbf{r} \times \rho_0 \mathbf{v} d\text{Vol} = -\frac{\mathbf{v}}{2c} \times \int \rho_0 \mathbf{r} d\text{Vol} = -\frac{\mathbf{v}}{2c} \times \mathbf{p}_0, \quad (3)$$

which differs from eq. (2) by a factor of 2.

Furthermore, support for this result appears to be given in probs. 6.21, 6.22 and 11.27 of [7].

However, we should recall that the origin of the first equality in (3) is a multipole expansion of the (quasistatic) vector potential of a current distribution. See, for example, sec. 5.6 of [7]. That form depends on the current density  $\mathbf{J}$  having zero divergence. In general,

$$\nabla \cdot \mathbf{J} = -\frac{\partial \rho}{\partial t}, \quad (4)$$

where  $\rho$  is the electric charge density.<sup>3</sup> The charge density of a moving electric dipole is time dependent, such that  $\nabla \cdot \mathbf{J} \neq 0$ , and we cannot expect the analysis of eq. (3) to be valid.<sup>4</sup>

## 2.2 The Fields of a Moving Electric Dipole

The literature on calculations of the fields of a moving, point electric dipole is very extensive, including [8, 9, 10, 11, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25]. Most of these works emphasize the radiation fields of an ultrarelativistic dipole. Here, we consider only the low-velocity limit of an electric dipole, of strength  $\mathbf{p}_0$  in its rest frame, that is at the origin at time  $t = 0$  with velocity  $\mathbf{v}$ . We work in the quasistatic approximation, in which the potentials are the same in the Coulomb and Lorenz gauges.

### 2.2.1 Multipole Expansions of the Potentials

Following the spirit of [6], and secs. 4.1 and 5.6 of [7], we consider multipole expansions of the scalar potential  $V$  and the vector potential  $\mathbf{A}$  at time  $t = 0$ , when the moving electric dipole is at the origin. Then, the (quasistatic) scalar potential is

$$V(\mathbf{r}, t = 0) = \frac{\mathbf{p}_0 \cdot \mathbf{r}}{r^3}, \quad (5)$$

as the charge distribution has only a dipole moment at this time. The (quasistatic) vector potential has the expansion

$$\begin{aligned} A_i(\mathbf{r}, t = 0) &= \frac{1}{cr} \int J_i(\mathbf{r}', t = 0) d\text{Vol}' + \frac{\mathbf{r}}{cr^3} \cdot \int \mathbf{r}' J_i(\mathbf{r}', t = 0) d\text{Vol}' + \dots \\ &= \frac{\mathbf{v}}{cr} \int \rho_0(\mathbf{r}', t = 0) d\text{Vol}' + \frac{v_i \mathbf{r}}{cr^3} \cdot \int \mathbf{r}' \rho_0(\mathbf{r}', t = 0) d\text{Vol}' + \dots \\ &= \frac{v_i \mathbf{r} \cdot \mathbf{p}_0}{cr^3}, \end{aligned} \quad (6)$$

---

<sup>3</sup>For moving media,  $\nabla \cdot \mathbf{J} = -\partial \rho / \partial t - (\mathbf{v} \cdot \nabla) \rho = -d\rho / dt$ .

<sup>4</sup>Thanks to Grigory Vekstein for pointing this out.

where  $\rho_0$  is the electric charge distribution of the electric dipole in its rest frame,  $\int \rho_0(\mathbf{r}) d\text{Vol} = 0$ ,  $\int \rho_0(\mathbf{r}) \mathbf{r} d\text{Vol} = \mathbf{p}_0$ , and  $\mathbf{J} = \rho_0 \mathbf{v}$  is the current density of the moving electric dipole in the lab frame. It is perhaps not evident that all higher-order terms vanish in eq. (6), but this will be confirmed in sec. 2.2.2. Then,

$$\mathbf{A}(\mathbf{r}, t = 0) = \frac{(\mathbf{p}_0 \cdot \mathbf{r})\mathbf{v}}{cr^3} = \left(-\frac{\mathbf{v}}{c} \times \mathbf{p}_0\right) \times \frac{\mathbf{r}}{r^3} + \frac{(\mathbf{r} \cdot \mathbf{v})\mathbf{p}_0}{cr^3} \equiv \mathbf{A}_m + \mathbf{A}_p, \quad (7)$$

where

$$\mathbf{A}_m = \frac{\mathbf{m} \times \mathbf{r}}{r^3} = \frac{(\mathbf{p}_0 \cdot \mathbf{r})\mathbf{v} - (\mathbf{r} \cdot \mathbf{v})\mathbf{p}_0}{cr^3} \quad \text{with} \quad \mathbf{m} = -\frac{\mathbf{v}}{c} \times \mathbf{p}_0, \quad (8)$$

which is the vector potential expected from eq. (2) for a moving electric dipole moment (with no intrinsic magnetic moment), and

$$\mathbf{A}_p = \frac{(\mathbf{r} \cdot \mathbf{v})\mathbf{p}_0}{cr^3}. \quad (9)$$

It is something of a convention to say that  $\mathbf{A}_m$  is the vector potential of the magnetic moment of the moving electric dipole in that the term  $\mathbf{A}_p$  is also a vector potential with  $1/r^2$  dependence as associated with dipole potentials. The decomposition  $\mathbf{A} = \mathbf{A}_m + \mathbf{A}_p$  is of possible mathematical convenience but has no well-defined physical significance. For example, we could also write

$$\mathbf{A} = \frac{(\mathbf{p}_0 \cdot \mathbf{r})\mathbf{v}}{cr^3} = \frac{(\mathbf{p}_0 \cdot \mathbf{r})\mathbf{v} - (\mathbf{r} \cdot \mathbf{v})\mathbf{p}_0}{2cr^3} + \frac{(\mathbf{p}_0 \cdot \mathbf{r})\mathbf{v} + (\mathbf{r} \cdot \mathbf{v})\mathbf{p}_0}{2cr^3} = \mathbf{A}_a + \mathbf{A}_s, \quad (10)$$

where

$$\mathbf{A}_a = \frac{(\mathbf{p}_0 \cdot \mathbf{r})\mathbf{v} - (\mathbf{r} \cdot \mathbf{v})\mathbf{p}_0}{2cr^3} = \frac{1}{2}\mathbf{A}_m, \quad \mathbf{A}_s = \frac{(\mathbf{p}_0 \cdot \mathbf{r})\mathbf{v} + (\mathbf{r} \cdot \mathbf{v})\mathbf{p}_0}{2cr^3}, \quad (11)$$

are the antisymmetric and symmetric combinations of the terms  $(\mathbf{p}_0 \cdot \mathbf{r})\mathbf{v}/2cr^3$  and  $(\mathbf{r} \cdot \mathbf{v})\mathbf{p}_0/2cr^3$ . The decomposition (10) is advocated in probs. 6.21 and 6.22 of [7], but should not be construed as implying that the magnetic moment of the moving electric dipole is  $\mathbf{A}_m/2$ , just as the decomposition (7) does not imply that the magnetic moment of the moving electric dipole is  $\mathbf{A}_m$ .<sup>5</sup>

It would seem most correct simply to say that  $1/r^2$  term in the vector potential of the moving electric dipole is  $(\mathbf{p}_0 \cdot \mathbf{r})\mathbf{v}/cr^3$  without supposing this to be the sum of two effects of different physical character. Nonetheless, we can be led to the decompositions (7) and (10) by other arguments, each compelling in its way, as considered in secs. 2.2.3 and 2.2.4.

## 2.2.2 Fields and Potentials via Lorentz Transformations

Before going further, it is useful to note the most direct method of obtaining the potentials and fields of a moving electric dipole is via a Lorentz transformation from its rest frame, which has velocity  $\mathbf{v}$  with  $v \ll c$  with respect to the lab frame.

---

<sup>5</sup>That Jackson was aware of both decompositions (7) and (10) is indicated in a letter of Mar. 20, 1990 on this theme [26], and somewhat indirectly in prob. 11.28 of [7]. *V.H. thanks J.D.J. for a copy of this letter.*

The potentials in the rest frame of the electric dipole, assumed to be at the origin, are

$$V^* = \frac{\mathbf{p}_0 \cdot \mathbf{r}^*}{r^{*3}}, \quad \mathbf{A}^* = 0, \quad (12)$$

where quantities in the rest frame are denoted with the superscript  $*$ . The Lorentz transformation of the 4-vectors  $(ct, \mathbf{r})$  and  $(V, \mathbf{A})$  to the lab frame at time  $t = 0$  yield, for  $v \ll c$ , where  $\gamma = 1/\sqrt{1 - v^2/c^2} \approx 1$ ,

$$\mathbf{r}_{\parallel}^* = \gamma(\mathbf{r}_{\parallel} - \mathbf{v}t) \approx \mathbf{r}_{\parallel}, \quad \mathbf{r}_{\perp}^* = \mathbf{r}_{\perp}, \quad i.e., \quad \mathbf{r}^* \approx \mathbf{r}, \quad (13)$$

$$V = \gamma(V^* + \mathbf{A}^* \cdot \mathbf{v}/c) \approx V^* = \frac{\mathbf{p}_0 \cdot \mathbf{r}^*}{r^{*3}} \approx \frac{\mathbf{p}_0 \cdot \mathbf{r}}{r^3}, \quad (14)$$

$$\mathbf{A} = \gamma(\mathbf{A}^* + V^* \mathbf{v}/c) \approx \frac{(\mathbf{p}_0 \cdot \mathbf{r}) \mathbf{v}}{cr^3}. \quad (15)$$

Hence, the multipole expansion (7) of the vector potential  $\mathbf{A}$  at order  $1/r^2$  is actually the “exact” result in the low-velocity approximation.

While the fields  $\mathbf{E}$  and  $\mathbf{B}$  can now be calculated from the potentials according to

$$\mathbf{E} = -\nabla V - \frac{1}{c} \frac{\partial \mathbf{A}}{\partial t}, \quad \mathbf{B} = \nabla \times \mathbf{A}, \quad (16)$$

it is more straightforward to deduce the fields  $\mathbf{E}$  and  $\mathbf{B}$  as the Lorentz transforms of the fields in the rest frame of the electric dipole,

$$\mathbf{E}^* = \frac{3(\mathbf{p}_0 \cdot \mathbf{r}^*) \mathbf{r}^*}{r^{*5}} - \frac{\mathbf{p}_0}{r^{*3}}, \quad \mathbf{B}^* = 0. \quad (17)$$

Then,<sup>6</sup> at time  $t = 0$  when  $\mathbf{r}^* \approx \mathbf{r}$ ,

$$\mathbf{E} = \gamma \left( \mathbf{E}^* - \frac{\mathbf{v}}{c} \times \mathbf{B}^* \right) - \frac{\gamma^2}{\gamma + 1} \frac{(\mathbf{E}^* \cdot \mathbf{v}) \mathbf{v}}{c^2} \approx \mathbf{E}^* = \frac{3(\mathbf{p}_0 \cdot \mathbf{r}) \mathbf{r}}{r^5} - \frac{\mathbf{p}_0}{r^3}, \quad (18)$$

$$\begin{aligned} \mathbf{B} &= \gamma \left( \mathbf{B}^* + \frac{\mathbf{v}}{c} \times \mathbf{E}^* \right) - \frac{\gamma^2}{\gamma + 1} \frac{(\mathbf{B}^* \cdot \mathbf{v}) \mathbf{v}}{c^2} \approx \frac{\mathbf{v}}{c} \times \mathbf{E}^* \approx \frac{\mathbf{v}}{c} \times \mathbf{E} \\ &= \frac{\mathbf{v}}{c} \times \left( \frac{3(\mathbf{p}_0 \cdot \mathbf{r}) \mathbf{r}}{r^5} - \frac{\mathbf{p}_0}{r^3} \right) = \mathbf{B}_m + \mathbf{B}_p = \mathbf{B}_a + \mathbf{B}_s, \end{aligned} \quad (19)$$

where the partial fields  $\mathbf{B}_m$ ,  $\mathbf{B}_p$ ,  $\mathbf{B}_a$  and  $\mathbf{B}_s$  will be displayed in secs. 2.2.3 and 2.2.4.

### 2.2.3 Potentials Deduced from Lab-Frame Charge and Current Densities

In its rest-frame the point electric dipole  $\mathbf{p}_0$  (located at the origin) has “free” polarization density

$$\mathbf{P}_{\text{free}}^* = \mathbf{p}_0 \delta^3 \mathbf{r}^*. \quad (20)$$

---

<sup>6</sup>See, for example, sec. 11.10 of [7].

The associated density of free charge is

$$\rho_{\text{free}}^* = -\nabla^* \cdot \mathbf{P}_{\text{free}}^* = -\mathbf{p}_0 \cdot \nabla^* \delta^3 \mathbf{r}^*. \quad (21)$$

In the lab frame the free-charge density is

$$\rho_{\text{free}} = \gamma \rho_{\text{free}}^* \approx \rho_{\text{free}}^* \approx -\mathbf{p}_0 \cdot \nabla \delta^3 \mathbf{r}, \quad (22)$$

at the instant when the dipole is at the origin, noting that  $\mathbf{r} \approx \mathbf{r}^*$  and  $\nabla^* = \nabla$  in the low-velocity approximation.

When the dipole is moving with a velocity  $\mathbf{v}$  in the lab frame it has a free-current density

$$\begin{aligned} \mathbf{J}_{\text{free}} &= \gamma \rho_{\text{free}}^* \mathbf{v} \approx \rho_{\text{free}}^* \mathbf{v} = -\mathbf{v}(\mathbf{p}_0 \cdot \nabla) \delta^3 \mathbf{r} \\ &= -(\mathbf{v} \cdot \nabla) \mathbf{p}_0 \delta^3(\mathbf{r}) - \mathbf{v}(\nabla \cdot \mathbf{p}_0 \delta^3 \mathbf{r}) + (\mathbf{v} \cdot \nabla) \mathbf{p}_0 \delta^3 \mathbf{r} \\ &= -(\mathbf{v} \cdot \nabla) \mathbf{p}_0 \delta^3 \mathbf{r} - \nabla \times (\mathbf{v} \times \mathbf{p}_0 \delta^3 \mathbf{r}) \\ &\equiv \mathbf{J}_p + \mathbf{J}_m, \end{aligned} \quad (23)$$

noting that the operator  $\nabla$  does not act on the constant vectors  $\mathbf{p}_0$  and  $\mathbf{v}$ . The current density  $\mathbf{J}_p$  can be thought of as an “electric-polarization current,”

$$\mathbf{J}_p = -(\mathbf{v} \cdot \nabla) \mathbf{p}_0 \delta^3 \mathbf{r} \equiv -(\mathbf{v} \cdot \nabla) \mathbf{P}_{\text{free}} = -\frac{d\mathbf{P}_{\text{free}}}{dt} \quad (24)$$

where

$$\mathbf{P}_{\text{free}} = \mathbf{p}_0 \delta^3 \mathbf{r}. \quad (25)$$

Likewise, the current density  $\mathbf{J}_m$  can be thought of as a “magnetic-polarization current,”

$$\mathbf{J}_m = -\nabla \times (\mathbf{v} \times \mathbf{p}_0 \delta^3 \mathbf{r}) = c \nabla \times \mathbf{m} \delta^3 \mathbf{r} \equiv c \nabla \times \mathbf{M}_{\text{free}}, \quad (26)$$

using the convention that  $\mathbf{m} = -\mathbf{v}/c \times \mathbf{p}_0$  of eq. (2), and

$$\mathbf{M}_{\text{free}} = \mathbf{m} \delta^3 \mathbf{r}. \quad (27)$$

Then the sum of the “free-polarization” currents (24) and (26) equals the “convection” current (23).

The (quasistatic) vector potentials associated with the free-polarization currents (24) and (26) when the electric dipole is at the origin in the lab frame are

$$\begin{aligned} \mathbf{A}_p(\mathbf{r}) &= \frac{1}{c} \int \frac{\mathbf{J}_p(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\text{Vol}' = -\frac{1}{c} \int \frac{(\mathbf{v} \cdot \nabla') \mathbf{p}_0 \delta^3 \mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|} d\text{Vol}' = -\frac{\mathbf{p}_0}{c} \int \frac{(\mathbf{v} \cdot \nabla') \delta^3 \mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|} d\text{Vol}' \\ &= \frac{\mathbf{p}_0}{c} \int \frac{[\mathbf{v} \cdot (\mathbf{r} - \mathbf{r}')] \delta^3 \mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|^3} d\text{Vol}' = \frac{(\mathbf{r} \cdot \mathbf{v}) \mathbf{p}_0}{cr^3}, \end{aligned} \quad (28)$$

as previously found in eq. (9), and

$$\begin{aligned} \mathbf{A}_m(\mathbf{r}) &= \frac{1}{c} \int \frac{\mathbf{J}_m(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\text{Vol}' = \int \frac{\nabla' \times \mathbf{m} \delta^3 \mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|} d\text{Vol}' = -\mathbf{m} \times \int \frac{\nabla' \delta^3 \mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|} d\text{Vol}' \\ &= \mathbf{m} \times \int \frac{(\mathbf{r} - \mathbf{r}') \delta^3 \mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|^3} d\text{Vol}' = \frac{\mathbf{m} \times \mathbf{r}}{r^3} = -\frac{(\mathbf{v} \times \mathbf{p}_0) \times \mathbf{r}}{cr^3} = \frac{(\mathbf{p}_0 \cdot \mathbf{r}) \mathbf{v} - (\mathbf{r} \cdot \mathbf{v}) \mathbf{p}_0}{cr^3}, \end{aligned} \quad (29)$$

as previously found in eq. (8).

The magnetic field (19) can now be written as

$$\mathbf{B} = \frac{\mathbf{v}}{c} \times \left( \frac{3(\mathbf{p}_0 \cdot \mathbf{r})\mathbf{r}}{r^5} - \frac{\mathbf{p}_0}{r^3} \right) = \mathbf{B}_m + \mathbf{B}_p. \quad (30)$$

where

$$\begin{aligned} \mathbf{B}_m &= \nabla \times \mathbf{A}_m = -\nabla \times \frac{(\mathbf{v} \times \mathbf{p}_0) \times \mathbf{r}}{cr^3} \\ &= -\frac{\mathbf{v} \times \mathbf{p}_0}{cr^3} \nabla \cdot \mathbf{r} + \mathbf{r} \left( \nabla \cdot \frac{\mathbf{v} \times \mathbf{p}_0}{cr^3} \right) - (\mathbf{r} \cdot \nabla) \frac{\mathbf{v} \times \mathbf{p}_0}{cr^3} + \left( \frac{\mathbf{v} \times \mathbf{p}_0}{cr^3} \cdot \nabla \right) \mathbf{r} \\ &= -\frac{3\mathbf{v} \times \mathbf{p}_0}{cr^3} - \frac{3[\mathbf{r} \cdot (\mathbf{v} \times \mathbf{p}_0)]\mathbf{r}}{cr^5} + \frac{3\mathbf{v} \times \mathbf{p}_0}{cr^3} + \frac{\mathbf{v} \times \mathbf{p}_0}{cr^3} \\ &= -\frac{3[(\mathbf{v} \times \mathbf{p}_0) \cdot \mathbf{r}]\mathbf{r}}{cr^5} + \frac{\mathbf{v} \times \mathbf{p}_0}{cr^3} \\ &= \nabla \times \frac{(\mathbf{p}_0 \cdot \mathbf{r})\mathbf{v} - (\mathbf{r} \cdot \mathbf{v})\mathbf{p}_0}{cr^3} \\ &= (\mathbf{p}_0 \cdot \mathbf{r})\nabla \times \frac{\mathbf{v}}{cr^3} - \frac{\mathbf{v}}{cr^3} \times \nabla(\mathbf{p}_0 \cdot \mathbf{r}) - (\mathbf{v} \cdot \mathbf{r})\nabla \times \frac{\mathbf{p}_0}{cr^3} + \frac{\mathbf{p}_0}{cr^3} \times \nabla(\mathbf{v} \cdot \mathbf{r}) \\ &= -\frac{3(\mathbf{p}_0 \cdot \mathbf{r})\mathbf{r} \times \mathbf{v}}{cr^5} - \frac{\mathbf{v} \times \mathbf{p}_0}{cr^3} + \frac{3(\mathbf{v} \cdot \mathbf{r})\mathbf{r} \times \mathbf{p}_0}{cr^5} + \frac{\mathbf{p}_0 \times \mathbf{v}}{cr^3} \\ &= \mathbf{v} \times \frac{3(\mathbf{p}_0 \cdot \mathbf{r})\mathbf{r}}{cr^5} - \frac{2\mathbf{v} \times \mathbf{p}_0}{cr^3} - \mathbf{p}_0 \times \frac{3(\mathbf{v} \cdot \mathbf{r})\mathbf{r}}{cr^5}, \end{aligned} \quad (31)$$

and

$$\begin{aligned} \mathbf{B}_p &= \nabla \times \mathbf{A}_p = \nabla \times \frac{(\mathbf{r} \cdot \mathbf{v})\mathbf{p}_0}{cr^3} = (\mathbf{r} \cdot \mathbf{v})\nabla \times \frac{\mathbf{p}_0}{cr^3} + \nabla(\mathbf{r} \cdot \mathbf{v}) \times \frac{\mathbf{p}_0}{cr^3} \\ &= -3(\mathbf{r} \cdot \mathbf{v})\mathbf{r} \times \frac{\mathbf{p}_0}{cr^5} + \mathbf{v} \times \frac{\mathbf{p}_0}{cr^3} = \mathbf{p}_0 \times \left( \frac{3(\mathbf{r} \cdot \mathbf{v})\mathbf{r}}{cr^5} - \frac{\mathbf{v}}{cr^3} \right) \\ &= \mathbf{v} \times \frac{\mathbf{p}_0}{cr^3} + \mathbf{p}_0 \times \frac{3(\mathbf{r} \cdot \mathbf{v})\mathbf{r}}{cr^5}. \end{aligned} \quad (32)$$

The consistency of this decomposition tends to reinforce the identification of eq. (26) as “the” magnetic moment of the moving electric dipole, even though the quantities  $\mathbf{A}_p$  and  $\mathbf{B}_p$  also have the  $1/r^2$  and  $1/r^3$  dependence, respectively, expected for dipole potentials and fields. However, alternative decompositions of the electric and magnetic lead to different interpretations. One such decomposition is considered in the following section.

#### 2.2.4 Jackson’s Argument

The preceding analyses have assumed for simplicity that the electric dipole is at the origin. Problems 6.21 and 6.22 of [7] encourage us to consider the situation when the electric dipole is not at the origin.

If the dipole is at position  $\mathbf{r}_0$ , most of the preceding results still hold with the substitutions  $\mathbf{r} \rightarrow \mathbf{r} - \mathbf{r}_0$  and  $r \rightarrow |\mathbf{r} - \mathbf{r}_0|$ . However, the multipole moments of charge and current densities depend on the choice of origin.

The total electric charge,  $Q = \int \rho d\text{Vol}$  is, of course, independent of the choice of origin. The electric dipole moment  $\mathbf{p} = \int \rho \mathbf{r} d\text{Vol}$  is independent of the choice of origin if the total charge  $Q$  is zero (as for an electric dipole moment). Hence, the electric dipole moment of a dipole charge distribution is independent of the choice of origin. However, the electric quadrupole moment of a dipole charge distribution depends on the choice of origin.

The magnetic moment of a current density that obeys  $\nabla \cdot \mathbf{J} = 0$  is independent of the choice of origin. In such cases  $\int \mathbf{J} d\text{vol} = 0$ , and we can write

$$\mathbf{m} = \frac{1}{c} \int \mathbf{r} \times \mathbf{J} d\text{Vol} \rightarrow \frac{1}{c} \int (\mathbf{r} - \mathbf{r}_0) \times \mathbf{J} d\text{Vol} = \mathbf{m} + \frac{\mathbf{r}_0}{c} \times \int \mathbf{J} d\text{Vol} = \mathbf{m}. \quad (33)$$

In the present example of a moving electric dipole the current density has nonzero divergence, and the magnetic dipole moment (and higher magnetic moments) depend on the choice of origin. This reinforces the sense of previous sections that it is difficult to identify “the” magnetic moment of a moving electric dipole.

Turning to the argument implied in probs. 6.21 and 6.22 of [7], it is suggested that the vector potential (15) be rewritten in the form (10) as the sum of antisymmetric and symmetric terms. The motivation here is not obvious in quasistatic examples, but this technique is of use when considering a Taylor expansion of the retarded vector potential of a time-harmonic current distribution  $\mathbf{J}(\mathbf{r}, t) = \mathbf{J}(\mathbf{r}) e^{-i\omega t}$ , at distances far from these currents,

$$\begin{aligned} \mathbf{A}(\mathbf{r}, t) &= \frac{1}{c} \int \frac{\mathbf{J}(\mathbf{r}', t' = t - R/c)}{R} d\text{Vol}' \approx \frac{e^{-i\omega t}}{cr} \int \mathbf{J}(\mathbf{r}') e^{ikR} d\text{Vol}' \\ &\approx \frac{e^{i(kr - \omega t)}}{cr} \int \mathbf{J}(\mathbf{r}') e^{-ik\hat{\mathbf{r}} \cdot \mathbf{r}'} d\text{Vol}' \\ &\approx \frac{e^{i(kr - \omega t)}}{cr} \left( \int \mathbf{J}(\mathbf{r}') d\text{Vol}' - ik \int \mathbf{J}(\mathbf{r}') (\hat{\mathbf{r}} \cdot \mathbf{r}') d\text{Vol}' + \dots \right), \end{aligned} \quad (34)$$

where  $R = |\mathbf{r} - \mathbf{r}'| \approx r - \hat{\mathbf{r}} \cdot \mathbf{r}'$ . As is well known (see, for example, secs. 9.2-3 of [7]), the first term corresponds to the vector potential of electric dipole radiation, and the second term is the sum of the vector potentials of magnetic dipole radiation and electric quadrupole radiation. The separation of the second term into the two types of radiation is accomplished by considering its symmetric and antisymmetric parts,

$$\begin{aligned} \mathbf{J}(\mathbf{r}') (\hat{\mathbf{r}} \cdot \mathbf{r}') &= \frac{(\hat{\mathbf{r}} \cdot \mathbf{r}') \mathbf{J}(\mathbf{r}') - (\hat{\mathbf{r}} \cdot \mathbf{J}(\mathbf{r}')) \mathbf{r}'}{2} + \frac{(\hat{\mathbf{r}} \cdot \mathbf{r}') \mathbf{J}' + (\hat{\mathbf{r}} \cdot \mathbf{J}(\mathbf{r}')) \mathbf{r}'}{2} \\ &= \frac{(\mathbf{r}' \times \mathbf{J}(\mathbf{r}')) \times \hat{\mathbf{r}}}{2} + \frac{(\hat{\mathbf{r}} \cdot \mathbf{r}') \mathbf{J}(\mathbf{r}') + (\hat{\mathbf{r}} \cdot \mathbf{J}(\mathbf{r}')) \mathbf{r}'}{2} \\ &= c\mathbf{M}(\mathbf{r}') \times \hat{\mathbf{r}} + \frac{(\hat{\mathbf{r}} \cdot \mathbf{r}') \mathbf{J}(\mathbf{r}') + (\hat{\mathbf{r}} \cdot \mathbf{J}(\mathbf{r}')) \mathbf{r}'}{2}, \end{aligned} \quad (35)$$

where  $\mathbf{M}(\mathbf{r}) = \mathbf{r} \times \mathbf{J}(\mathbf{r})/2c$  is the magnetization density associated with current density  $\mathbf{J}$ .

*The identification of  $\mathbf{r} \times \mathbf{J}/2c$  as a magnetization density seems justified if  $\mathbf{J}$  has zero divergence, which is not the case, in general, in radiation problems. It is noteworthy that [27] (sec. 71) discusses both magnetic moments and radiation for a collection of point charges, rather than for a current density  $\mathbf{J}$ , which deftly avoids the present ambiguities.*

For completeness, we record that the magnetic field associated with the symmetric part,  $\mathbf{A}_s$ , of the vector potential (10) is, at time  $t = 0$  when the moving electric dipole is at the origin,

$$\begin{aligned}
\mathbf{B}_s &= \nabla \times \mathbf{A}_s = \nabla \times \frac{(\mathbf{p}_0 \cdot \mathbf{r})\mathbf{v} + (\mathbf{r} \cdot \mathbf{v})\mathbf{p}_0}{2cr^3} \\
&= (\mathbf{p}_0 \cdot \mathbf{r})\nabla \times \frac{\mathbf{v}}{2cr^3} - \frac{\mathbf{v}}{2cr^3} \times \nabla(\mathbf{p}_0 \cdot \mathbf{r}) + (\mathbf{v} \cdot \mathbf{r})\nabla \times \frac{\mathbf{p}_0}{2cr^3} - \frac{\mathbf{p}_0}{2cr^3} \times \nabla(\mathbf{v} \cdot \mathbf{r}) \\
&= -\frac{3(\mathbf{p}_0 \cdot \mathbf{r})\mathbf{r} \times \mathbf{v}}{2cr^5} - \frac{\mathbf{v} \times \mathbf{p}_0}{2cr^3} - \frac{3(\mathbf{v} \cdot \mathbf{r})\mathbf{r} \times \mathbf{p}_0}{2cr^5} - \frac{\mathbf{p}_0 \times \mathbf{v}}{2r^3} \\
&= \mathbf{v} \times \frac{3(\mathbf{p}_0 \cdot \mathbf{r})\mathbf{r}}{2cr^5} + \mathbf{p}_0 \times \frac{3(\mathbf{v} \cdot \mathbf{r})\mathbf{r}}{2cr^5} = -\frac{3\mathbf{r} \times [(\mathbf{v} \cdot \mathbf{r})\mathbf{p}_0 + (\mathbf{p}_0 \cdot \mathbf{r})\mathbf{v}]}{2cr^5}. \tag{36}
\end{aligned}$$

When the moving dipole is at the origin the magnetic field varies purely as  $1/r^3$ , so the fourth Maxwell equation (for points away from the dipole itself),

$$\nabla \times \mathbf{B} = \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t}, \tag{37}$$

tells us that the electric field must include terms that vary as  $1/r^4$ , which are associated with the (time-dependent) electric quadrupole moment. However, eq. (37) should not be construed as implying that the changing electric quadrupole moment “creates” the magnetic field.

Since  $\mathbf{B}_m = 2\mathbf{B}_a$  has the form of a static dipole field, the curls of these fields are zero away from the dipole, and in that region we can write

$$\nabla \times \mathbf{B}_p = \nabla \times \mathbf{B}_s = \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t}. \tag{38}$$

This establishes a relation between  $\mathbf{B}_p$ ,  $\mathbf{B}_s$  and the electric quadrupole field of the moving electric dipole, but this should not be interpreted as a cause-and-effect relation.

It remains that the magnetic field of a moving electric dipole  $\mathbf{p}_0$  at order  $1/r^3$  is not entirely due to the “magnetic moment”  $-\mathbf{v}/c \times \mathbf{p}_0$ , so that it is somewhat delicate to use the terminology “magnetic moment” in this example.

### 3 Moving Magnetic Dipole

For completeness, we include a discussion of the case of a magnetic dipole  $\mathbf{m}_0$  that has velocity  $\mathbf{v}$  in the lab frame.

The potentials in the rest frame of magnetic dipole (where the magnetic moment is  $\mathbf{m}_0$ ), assuming the dipole to be at the origin, are

$$V^* = 0, \quad \mathbf{A}^* = \frac{\mathbf{m}_0 \times \mathbf{r}^*}{r^{*3}}, \tag{39}$$



where quantities in the rest frame are denoted with the superscript  $\star$ . The Lorentz transformation of the 4-vector  $(V, \mathbf{A})$  to the lab frame at time  $t = 0$  yield, for  $v \ll c$ , where  $\gamma = 1/\sqrt{1 - v^2/c^2} \approx 1$ , and  $\mathbf{r}^\star \approx \mathbf{r}$ , is

$$V = \gamma(V^\star + \mathbf{A}^\star \cdot \mathbf{v}/c) \approx \frac{\mathbf{m}_0 \times \mathbf{r} \cdot \mathbf{v}}{cr^3} = \frac{\mathbf{v}/c \times \mathbf{m}_0 \cdot \mathbf{r}}{r^3}, \quad (40)$$

$$\mathbf{A} = \gamma(\mathbf{A}^\star + V^\star \mathbf{v}/c) \approx \mathbf{A}^\star = \frac{\mathbf{m}_0 \times \mathbf{r}}{r^3}. \quad (41)$$

These potentials have a crisper interpretation than those, (14)-(15) for a moving electric dipole, in that we can say that a moving magnetic dipole has magnetic dipole moment  $\mathbf{m} = \mathbf{m}_0$  and electric dipole moment

$$\mathbf{p} = \frac{\mathbf{v}}{c} \times \mathbf{m}_0, \quad (42)$$

with respect to its instantaneous position.

The lab-frame fields  $\mathbf{E}$  and  $\mathbf{B}$  are the Lorentz transforms of the fields in the rest frame of the magnetic dipole,

$$\mathbf{E}^\star = 0, \quad \mathbf{B} = \frac{3(\mathbf{m}_0 \cdot \mathbf{r}^\star)\mathbf{r}^\star}{r^{\star 5}} - \frac{\mathbf{m}_0}{r^{\star 3}}. \quad (43)$$

Then,

$$\mathbf{E} \approx -\frac{\mathbf{v}}{c} \times \mathbf{B}^\star \approx -\frac{\mathbf{v}}{c} \times \mathbf{B} = -\frac{\mathbf{v}}{c} \times \left( \frac{3(\mathbf{m}_0 \cdot \mathbf{r})\mathbf{r}}{r^5} - \frac{\mathbf{m}_0}{r^3} \right), \quad (44)$$

$$\mathbf{B} \approx \mathbf{B}^\star = \frac{3(\mathbf{m}_0 \cdot \mathbf{r})\mathbf{r}}{r^5} - \frac{\mathbf{m}_0}{r^3}, \quad (45)$$

The electric field can also be written as

$$\mathbf{E} = \mathbf{E}_p + \mathbf{E}_m, \quad (46)$$

with

$$\mathbf{E}_p = -\nabla V = \frac{3(\mathbf{p} \cdot \mathbf{r})\mathbf{r}}{r^5} - \frac{\mathbf{p}}{r^3} = \frac{3(\mathbf{v} \times \mathbf{m}_0 \cdot \mathbf{r})\mathbf{r}}{cr^5} - \frac{\mathbf{v}}{c} \times \frac{\mathbf{m}_0}{r^3}, \quad (47)$$

and<sup>7</sup>

$$\mathbf{E}_m = -\frac{1}{c} \frac{\partial \mathbf{A}}{\partial t} = -\frac{1}{c} \frac{\partial}{\partial t} \frac{\mathbf{m}_0 \times \mathbf{r}}{r^3} = \mathbf{m}_0 \times \left( \frac{3(\mathbf{v} \cdot \mathbf{r})\mathbf{r}}{cr^5} - \frac{\mathbf{v}}{cr^3} \right), \quad (48)$$

where  $\mathbf{E}_p$  and  $\mathbf{E}_m$  can also be interpreted as the electric fields associated with the polarization and magnetization densities of the moving magnetic dipole, respectively.

---

<sup>7</sup>The form (48) was noted in prob. 12.4 of [28] and in prob. 11.29 of [7].

## References

- [1] V. Hnizdo, *Magnetic dipole moment of a moving electric dipole* (Jan. 4, 2012), <http://arxiv.org/abs/1201.0938>
- [2] H.A. Lorentz, *Alte und Neue Fragen der Physik*, Phys. Z. **11**, 1234 (1910), [http://puhep1.princeton.edu/~mcdonald/examples/EM/lorentz\\_pz\\_11\\_1234\\_10.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/lorentz_pz_11_1234_10.pdf)
- [3] I.M. Frank, *Izv. Akad. Nauk USSR, Ser. Fiz.* **6**, 3 (1942).
- [4] I.M. Frank, *Vavilov-Cherenkov radiation for electric and magnetic multipoles*, Sov. Phys. Usph. **27**, 772 (1984), [http://puhep1.princeton.edu/~mcdonald/examples/EM/frank\\_spu\\_27\\_772\\_84.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/frank_spu_27_772_84.pdf)
- [5] I.M. Frank, *On the moments of a magnetic dipole moving in a medium*, Sov. Phys. Usph. **32**, 456 (1989), [http://puhep1.princeton.edu/~mcdonald/examples/EM/frank\\_spu\\_32\\_456\\_89.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/frank_spu_32_456_89.pdf)
- [6] G.P. Fisher, *The Electric Dipole Moment of a Moving Magnetic Dipole*, Am. J. Phys. **39**, 1528 (1971), [http://puhep1.princeton.edu/~mcdonald/examples/EM/fisher\\_ajp\\_39\\_1528\\_71.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/fisher_ajp_39_1528_71.pdf)
- [7] J.D. Jackson, *Classical Electrodynamics*, 3rd ed. (Wiley, New York, 1999), [http://puhep1.princeton.edu/~mcdonald/examples/EM/jackson\\_ce3\\_pages.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/jackson_ce3_pages.pdf)
- [8] J.R. Ellis, *The fields of an arbitrarily moving dipole*, Proc. Camb. Phil. Soc. **59**, 759 (1963), [http://puhep1.princeton.edu/~mcdonald/examples/EM/ellis\\_pcps\\_59\\_759\\_63.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/ellis_pcps_59_759_63.pdf)
- [9] G.N. Ward, *On the integration of Maxwell's equations, and charge and dipole fields*, Proc. Roy. Soc. London A **279**, 562 (1964), [http://puhep1.princeton.edu/~mcdonald/examples/EM/ward\\_prsla\\_279\\_562\\_64.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/ward_prsla_279_562_64.pdf)
- [10] G.N. Ward, *The electromagnetic fields of moving dipoles*, Proc. Camb. Phil. Soc. **61**, 547 (1965), [http://puhep1.princeton.edu/~mcdonald/examples/EM/ward\\_pcps\\_61\\_547\\_65.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/ward_pcps_61_547_65.pdf)
- [11] H. Mott, G.B. Hoadley and W.R. Daw, *The Fields of a Moving Electric Dipole*, Proc. IEEE **53**, 407 (1965), [http://puhep1.princeton.edu/~mcdonald/examples/EM/mott\\_pieee\\_53\\_407\\_65.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/mott_pieee_53_407_65.pdf)
- [12] H. Mott, G.B. Hoadley and W.R. Daw, *The Near Fields of a Moving Electric Dipole*, Proc. IEEE **53**, 1262 (1965), [http://puhep1.princeton.edu/~mcdonald/examples/EM/mott\\_pieee\\_53\\_1262\\_65.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/mott_pieee_53_1262_65.pdf)
- [13] J.R. Ellis, *Electromagnetic Fields of Moving Dipoles and Multipoles*, J. Math. Phys. **7**, 1185 (1966), [http://puhep1.princeton.edu/~mcdonald/examples/EM/ellis\\_jmp\\_7\\_1185\\_66.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/ellis_jmp_7_1185_66.pdf)
- [14] H. Mott, G.B. Hoadley and W.R. Daw, *Radiation from the Moving Electric Dipole*, Proc. IEEE **55**, 92 (1967), [http://puhep1.princeton.edu/~mcdonald/examples/EM/mott\\_pieee\\_55\\_92\\_67.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/mott_pieee_55_92_67.pdf)

- [15] H. Mott, G.B. Hoadley and W.R. Daw, *The Fields of an Electric Dipole in Motion*, J. Franklin Inst. **283**, 91 (1967),  
[http://puhep1.princeton.edu/~mcdonald/examples/EM/mott\\_jfi\\_283\\_91\\_67.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/mott_jfi_283_91_67.pdf)
- [16] J.J. Monaghan, *The Heaviside-Feynman expression for the fields of an accelerated dipole*, J. Phys. A **1**, 112 (1969),  
[http://puhep1.princeton.edu/~mcdonald/examples/EM/monaghan\\_jpa\\_1\\_112\\_68.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/monaghan_jpa_1_112_68.pdf)
- [17] J.R. Ellis, *Force and couple exerted on a moving electromagnetic dipole*, J. Phys. A **3**, 251 (1970), [http://puhep1.princeton.edu/~mcdonald/examples/EM/ellis\\_jpa\\_3\\_251\\_70.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/ellis_jpa_3_251_70.pdf)
- [18] J. Heras, *Radiation fields of a dipole in arbitrary motion*, Am. J. Phys. **62**, 1109 (1984),  
[http://puhep1.princeton.edu/~mcdonald/examples/EM/heras\\_ajp\\_62\\_1109\\_94.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/heras_ajp_62_1109_94.pdf)
- [19] D. Schieber, *Radiation of a Receding Electric Dipole*, Electromagnetics **6**, 99 (1986),  
[http://puhep1.princeton.edu/~mcdonald/examples/EM/shieber\\_electromagnetics\\_6\\_99\\_86.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/shieber_electromagnetics_6_99_86.pdf)
- [20] J. Heras, *New approach to the classical radiation fields of moving dipoles*, Phys. Lett. A **237**, 343 (1998),  
[http://puhep1.princeton.edu/~mcdonald/examples/EM/heras\\_pl\\_a237\\_343\\_98.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/heras_pl_a237_343_98.pdf)
- [21] J. Heras, *Explicit expressions for the electric and magnetic fields of a moving magnetic dipole*, Phys. Rev. E **58**, 5047 (1998),  
[http://puhep1.princeton.edu/~mcdonald/examples/EM/heras\\_pre\\_58\\_5047\\_98.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/heras_pre_58_5047_98.pdf)
- [22] G.N. Afanasiev and Yu P. Stepanovksy, *Electromagnetic Fields of Electric, Magnetic and Toroidal Dipoles Moving in Medium*, Phys. Scripta **61**, 704 (2000),  
[http://puhep1.princeton.edu/~mcdonald/examples/EM/afanasiev\\_ps\\_61\\_704\\_00.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/afanasiev_ps_61_704_00.pdf)
- [23] E.A. Power and T. Thurinamachandran, *Electromagnetic fields of a moving electric dipole*, Proc. Roy. Soc. London A **457**, 2757 (2001),  
[http://puhep1.princeton.edu/~mcdonald/examples/EM/power\\_prsla\\_457\\_2757\\_01.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/power_prsla_457_2757_01.pdf)
- [24] E.A. Power and T. Thurinamachandran, *Radiative Properties of a moving electric dipole*, Proc. Roy. Soc. London A **457**, 2779 (2001),  
[http://puhep1.princeton.edu/~mcdonald/examples/EM/power\\_prsla\\_457\\_2779\\_01.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/power_prsla_457_2779_01.pdf)
- [25] G.S. Smith, *Visualizing special relativity: the field of an electric dipole moving at relativistic speed*, Eur. J. Phys. **32**, 695 (2011),  
[http://puhep1.princeton.edu/~mcdonald/examples/EM/smith\\_ejp\\_32\\_695\\_11.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/smith_ejp_32_695_11.pdf)
- [26] J.D. Jackson, Letter to W.A. Newcomb (Mar. 28, 1990),  
[http://puhep1.princeton.edu/~mcdonald/examples/EM/jackson\\_032890.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/jackson_032890.pdf)
- [27] L.D. Landau and E.M. Lifshitz, *The Classical Theory of Fields*, 4th ed. (Pergamon, 1975), [http://puhep1.princeton.edu/~mcdonald/examples/EM/landau\\_cft\\_sec71.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/landau_cft_sec71.pdf)
- [28] E.J. Konopinski, *Electromagnetic Fields and Relativistic Particles* (McGraw-Hill, 1981),  
[http://puhep1.princeton.edu/~mcdonald/examples/EM/konopinski\\_prob12-3-4\\_81.pdf](http://puhep1.princeton.edu/~mcdonald/examples/EM/konopinski_prob12-3-4_81.pdf)