

The Evening Sky Map

FREE* EACH MONTH FOR YOU TO EXPLORE, LEARN & ENJOY THE NIGHT SKY

WWW.SKYMAPS.COM

NORTHERN HEMISPHERE
JULY 2008

Sky Calendar – July 2008

- Mars 0.67° from Regulus** at 15h UT (50° from Sun, evening sky). Mags. +1.6 and +1.3. Closest a planet gets to a Mag. 1 star all year.
- Mercury at greatest elongation**, 22° west from Sun (morning sky) at 18h UT. Mag. +0.5. Look low in the east-northeast about 30 minutes before sunrise.
- Moon at perigee** (closest to Earth) at 21h UT (359,513 km; 33.2').
- New Moon** at 2:19 UT. Start of lunation 1058.
- Earth at Aphelion** (farthest from Sun) at 8h UT. The Sun-Earth distance is 1.01675 a.u. or about 152.1 million km.
- Moon near Beehive cluster (M44)** at 21h UT (24° from Sun, evening sky).
- Moon near Regulus** at 11h UT (evening sky).
- Moon near Mars** at 16h UT (evening sky). Mag. +1.7.
- Moon near Saturn** at 20h UT (evening sky). Mag. +0.8.
- Jupiter at opposition** at 8h UT (mag. -2.7). The best time to observe the largest planet in the solar system.
- First Quarter Moon** at 4:35 UT.
- Moon near Spica** at 16h UT (evening sky).
- Mars 0.64° from Saturn** at 16h UT (47° from Sun, evening sky). Mags. +1.7 and +0.8.
- Moon at apogee** (farthest from Earth) at 4h UT (distance 405,452 km; angular size 29.5').
- Moon very near Antares** at 12h UT (evening sky). Occultation visible from Southern Australia and N.Z.
- Moon near Jupiter** at 14h UT (midnight sky). Mag. -2.7.
- Full Moon** at 7:59 UT. The full Moon of July is called the "Thunder Moon" or "Hay Moon".
- Last Quarter Moon** at 18:42 UT.
- Moon near the Pleiades** at 17h UT (morning sky).
- Mercury at superior conjunction** at 20h UT (not visible). Passes into the evening sky.
- Moon at perigee** (closest to Earth) at 23h UT (363,883 km; 32.8').


All times in Universal Time (UT). (USA Eastern Summer Time = UT - 4 hours.)



SAVE ON RECOMMENDED PRODUCTS • <http://Skymaps.com/store>

- STAR ATLAS & PLANISPHERES
- STAR CHARTS & ASTRO POSTERS
- BOOKS FOR SKY WATCHERS
- TELESCOPES & BINOCULARS

All sales support the production and free distribution of The Evening Sky Map.



SKY MAP SHOWS HOW THE NIGHT SKY LOOKS

EARLY JUL 10 PM
LATE JUL 9 PM

(Add 1 Hour For Daylight Saving)
SKY MAP DRAWN FOR
A LATITUDE OF 40°
NORTH AND IS
SUITABLE FOR
LATITUDES UP
TO 15° NORTH
OR SOUTH
OF THIS

Symbols

- Galaxy ☁
- Double Star ●●
- Variable Star ⊙
- Diffuse Nebula □
- Planetary Nebula ⊕
- Open Star Cluster ☆
- Globular Star Cluster ⊕

Star Magnitudes ● ● ● ● ● ● ● ● ● ●
-1 0 1 2 3 4

Copyright © 2000–2008 Kym Thalassoudis. All Rights Reserved.

* TERMS OF USE: FREE FOR NON-COMMERCIAL EDUCATIONAL USE. ASTRONOMY EDUCATION GROUPS MAY FREELY DISTRIBUTE PRINTED HANDOUTS. FULL DETAILS AT <http://Skymaps.com/terms.html>

Low in the southern sky lies the bright and easy to recognize constellation known as Scorpius (The Scorpion).

INSTRUCTIONS: THE SKY MAP SHOWS THE ENTIRE NIGHT SKY FROM HORIZON-TO-HORIZON AS IT APPEARS ON CERTAIN DATES AND TIMES. THE CENTER OF THE MAP IS THE CENTER OF GREEK MYTHOLOGY.

About the Celestial Objects

Listed on this page are several of the brighter, more interesting celestial objects visible in the evening sky this month (refer to the monthly sky map). The objects are grouped into three categories. Those that can be easily seen with the naked eye (that is, without optical aid), those easily seen with binoculars, and those requiring a telescope to be appreciated. **Note, all of the objects (except single stars) will appear more impressive when viewed through a telescope or very large binoculars.** They are grouped in this way to highlight objects that can be seen using the optical equipment that may be available to the star gazer.

Tips for Observing the Night Sky

When observing the night sky, and in particular deep-sky objects such as star clusters, nebulae, and galaxies, it's always best to observe from a dark location. Avoid direct light from street lights and other sources. If possible observe from a dark location away from the light pollution that surrounds many of today's large cities.

You will see more stars after your eyes adapt to the darkness—usually about 10 to 20 minutes after you go outside. Also, if you need to use a torch to view the sky map, cover the light bulb with red cellophane. This will preserve your dark vision.

Finally, even though the Moon is one of the most stunning objects to view through a telescope, its light is so bright that it brightens the sky and makes many of the fainter objects very difficult to see. So try to observe the evening sky on moonless nights around either New Moon or Last Quarter.

Astronomical Glossary

Conjunction – An alignment of two celestial bodies such that they present the least angular separation as viewed from Earth.

Constellation – A defined area of the sky containing a star pattern.

Diffuse Nebula – A cloud of gas illuminated by nearby stars.

Double Star – Two stars that appear close to each other in the sky; either linked by gravity so that they orbit each other (binary star) or lying at different distances from Earth (optical double). Apparent separation of stars is given in seconds of arc (").

Ecliptic – The path of the Sun's center on the celestial sphere as seen from Earth.

Elongation – The angular separation of two celestial bodies. For Mercury and Venus the greatest elongation occurs when they are at their most angular distance from the Sun as viewed from Earth.

Galaxy – A mass of up to several billion stars held together by gravity.

Globular Star Cluster – A ball-shaped group of several thousand old stars.

Light Year (ly) – The distance a beam of light travels at 300,000 km/sec in one year.

Magnitude – The brightness of a celestial object as it appears in the sky.

Open Star Cluster – A group of tens or hundreds of relatively young stars.

Opposition – When a celestial body is opposite the Sun in the sky.

Planetary Nebula – The remnants of a shell of gas blown off by a star.

Universal Time (UT) – A time system used by astronomers. Also known as Greenwich Mean Time. USA Eastern Standard Time (for example, New York) is 5 hours behind UT.

Variable Star – A star that changes brightness over a period of time.

Easily Seen with the Naked Eye

| | | |
|------------|-----|----------------------------------------------------------------------------------------------|
| Altair | Aql | • Brightest star in Aquila. Name means "the flying eagle". Dist=16.7 ly. |
| Arcturus | Boo | • Orange, giant K star. Name means "bear watcher". Dist=36.7 ly. |
| δ Cephei | Cep | • Cepheid prototype. Mag varies between 3.5 & 4.4 over 5.366 days. Mag 6 companion. |
| Deneb | Cyg | • Brightest star in Cygnus. One of the greatest known supergiants. Dist=1,400±200 ly. |
| α Herculis | Her | • Semi-regular variable. Magnitude varies between 3.1 & 3.9 over 90 days. Mag 5.4 companion. |
| Vega | Lyr | • The 5th brightest star in the sky. A blue-white star. Dist=25.0 ly. |
| Antares | Sco | • Red, supergiant star. Name means "rival of Mars". Dist=135.9 ly. |
| Polaris | UMi | • The North Pole Star. A telescope reveals an unrelated mag 8 companion star. Dist=433 ly. |
| Spica | Vir | • Latin name means "ear of wheat" and shown held in Virgo's left hand. Dist=250 ly. |

Easily Seen with Binoculars

| | | |
|---------------|-----|----------------------------------------------------------------------------------------------|
| η Aquilae | Aql | • Bright Cepheid variable. Mag varies between 3.6 & 4.5 over 7.166 days. Dist=1,200 ly. |
| M3 | CVn | • Easy to find in binoculars. Might be glimpsed with the naked eye. |
| μ Cephei | Cep | • Herschel's Garnet Star. One of the reddest stars. Mag 3.4 to 5.1 over 730 days. |
| Mel 111 | Com | • Coma Berenices. 80 mag 5-6 stars in 5 deg. Dist=283 ly. Age=400 million years. |
| χ Cygni | Cyg | • Long period pulsating red giant. Magnitude varies between 3.3 & 14.2 over 407 days. |
| M39 | Cyg | • May be visible to the naked eye under good conditions. Dist=900 ly. |
| ν Draconis | Dra | • Wide pair of white stars. One of the finest binocular pairs in the sky. Dist=100 ly. |
| M13 | Her | • Best globular in northern skies. Discovered by Halley in 1714. Dist=23,000 ly. |
| M92 | Her | • Fainter and smaller than M13. Use a telescope to resolve its stars. |
| ε Lyrae | Lyr | • Famous Double Double. Binoculars show a double star. High power reveals each a double. |
| R Lyrae | Lyr | • Semi-regular variable. Magnitude varies between 3.9 & 5.0 over 46.0 days. |
| M12 | Oph | • Close to the brighter M10. Dist=18,000 ly. |
| M10 | Oph | • 3 degrees from the fainter M12. Both may be glimpsed in binoculars. Dist=14,000 ly. |
| IC 4665 | Oph | • Large, scattered open cluster. Visible with binoculars. |
| 6633 | Oph | • Scattered open cluster. Visible with binoculars. |
| M15 | Peg | • Only globular known to contain a planetary nebula (Mag 14, d=1"). Dist=30,000 ly. |
| M8 | Sgr | • Lagoon Nebula. Bright nebula bisected by a dark lane. Dist=5,200 ly. |
| M25 | Sgr | • Bright cluster located about 6 deg N of "teapot's" lid. Dist=1,900 ly. |
| M22 | Sgr | • A spectacular globular star cluster. Telescope will show stars. Dist=10,000 ly. |
| M4 | Sco | • A close globular. May just be visible without optical aid. Dist=7,000 ly. |
| M6 | Sco | • Butterfly Cluster. 30+ stars in 7x binoculars. Dist=1,960 ly. |
| M7 | Sco | • Superb open cluster. Visible to the naked eye. Age=260 million years. Dist=780 ly. |
| M5 | Ser | • Fine globular star cluster. Telescope will reveal individual stars. Dist=25,000 ly. |
| Mizar & Alcor | UMa | • Good eyesight or binoculars reveals 2 stars. Not a binary. Mizar has a mag 4 companion. |
| Cr 399 | Vul | • Coathanger asterism or "Brocchi's Cluster". Not a true star cluster. Dist=218 to 1,140 ly. |

Telescopic Objects

| | | |
|------------|-----|-----------------------------------------------------------------------------------------------|
| 7009 | Aqr | • Saturn Nebula. Requires 8-inch telescope to see Saturn-like appendages. |
| ε Boötis | Boo | • Red giant star (mag 2.5) with a blue-green mag 4.9 companion. Sep=2.8". Difficult to split. |
| M94 | CVn | • Compact nearly face-on spiral galaxy. Dist=15 million ly. |
| M51 | CVn | • Whirlpool Galaxy. First recognised to have spiral structure. Dist=25 million ly. |
| M64 | Com | • Black-Eye Galaxy. Discovered by J.E. Bode in 1775 - "a small, nebulous star". |
| Albireo | Cyg | • Beautiful double star. Contrasting colours of orange and blue-green. Sep=34.4". |
| 61 Cygni | Cyg | • Attractive double star. Mags 5.2 & 6.1 orange dwarfs. Dist=11.4 ly. Sep=28.4". |
| γ Delphini | Del | • Appear yellow & white. Mags 4.3 & 5.2. Dist=100 ly. Struve 2725 double in same field. |
| β Lyrae | Lyr | • Eclipsing binary. Mag varies between 3.3 & 4.3 over 12.940 days. Fainter mag 7.2 blue star. |
| M57 | Lyr | • Ring Nebula. Magnificent object. Smoke-ring shape. Dist=4,100 ly. |
| M23 | Sgr | • Elongated star cluster. Telescope required to show stars. Dist=2,100 ly. |
| M20 | Sgr | • Trifid Nebula. A telescope shows 3 dust lanes trisecting nebula. Dist=5,200 ly. |
| M21 | Sgr | • A fine and impressive cluster. Dist=4,200 ly. |
| M17 | Sgr | • Omega Nebula. Contains the star cluster NGC 6618. Dist=4,900 ly. |
| M11 | Sct | • Wild Duck Cluster. Resembles a globular through binoculars. V-shaped. Dist=5,600 ly. |
| M16 | Ser | • Eagle Nebula. Requires a telescope of large aperture. Dist=8,150 ly. |
| M81 | UMa | • Beautiful spiral galaxy visible with binoculars. Easy to see in a telescope. |
| M82 | UMa | • Close to M81 but much fainter and smaller. |
| M27 | Vul | • Dumbbell Nebula. Large, twin-lobed shape. Most spectacular planetary. Dist=975 ly. |